

# The Qingdao Twin Registry: A Status Update

Haiping Duan,<sup>1\*</sup> Feng Ning,<sup>2\*</sup> Dongfeng Zhang,<sup>3</sup> Shaojie Wang,<sup>1</sup> Dong Zhang,<sup>4</sup> Qihua Tan,<sup>5,6</sup> Xiaocao Tian,<sup>1,7</sup> and Zengchang Pang<sup>1</sup>

<sup>1</sup>*Qingdao Centers for Disease Control and Prevention, Qingdao, China*

<sup>2</sup>*Department of Public Health, Hjelt Institute, University of Helsinki, Helsinki, Finland*

<sup>3</sup>*Department of Public Health, Qingdao University Medical College, Qingdao, China*

<sup>4</sup>*Jiaonan Centers for Disease Control and Prevention, Qingdao, China*

<sup>5</sup>*Epidemiology, Institute of Public Health, University of Southern Denmark, Odense, Denmark*

<sup>6</sup>*Department of Clinical Genetics, Odense University Hospital, Odense, Denmark*

<sup>7</sup>*Department of Public Health, University of Shandong, Jinan, China*

\*These authors contributed equally.

In 1998, the Qingdao Twin Registry was initiated as the main part of the Chinese National Twin Registry. By 2005, a total of 10,655 twin pairs had been recruited. Since then new twin cohorts have been sampled, with one longitudinal cohort of adolescent twins selected to explore determinants of metabolic disorders and health behaviors during puberty and young adulthood. Adult twins have been sampled for studying heritability of multiple phenotypes associated with metabolic disorders. In addition, an elderly twin cohort has been recruited with a focus on genetic studies of aging-related phenotypes using twin modeling and genome-wide association analysis. Cross-cultural collaborative studies have been carried out between China, Denmark, Finland, and US cohorts. Ongoing data collection and analysis for the Qingdao Twin Registry will be discussed in this article.

■ **Keywords:** Chinese twins, registry, phenotypes, genome-wide association analysis, cohort study

During the past decades, non-communicable diseases have been the leading cause of mortality and morbidity worldwide. Developing countries emerged as epicenters of diabetes and other non-communicable diseases as a result of urbanization and nutritional transition (Chan et al., 2009; Wang et al., 2005; WHO, 1994). Non-communicable diseases account for an estimated 80% of annual deaths (Wang et al., 2005) and 70% of total disability-adjusted life-years in China ('China's major health challenge', 2011). Studying the etiology and causes of non-communicable diseases can have significant impacts on disease prevention and treatment.

The Qingdao Twin Registry (QTR), established by Qingdao Centers for Disease Control and Prevention (QDCDC) in 1998, is part of the Chinese National Twin Registry (Li et al., 2006). The purpose of the QTR was primarily to recruit twins born in the Qingdao region and to estimate genetic and environmental components in non-communicable diseases (Pang et al., 2006). A total of 10,655 twin pairs across all age groups were recruited until 2005 (Pang et al., 2006). The coverage of QTR has been extended and new cohorts were established during the past 5 years. Meanwhile, cross-culture collaborative studies between China, Denmark, Finland, and US cohorts have been

carried out based on the QTR. This paper provides a detailed status update of the QTR.

## Twins Recruitment

Since 2000, the QTR recruited twins through medical records, school, and media coverage (Pang et al., 2003, 2006). Newborn twins were identified and recruited to QTR through the Immunization Registry for Newborns, which covers 95% of all newborns in Qingdao each year. Children and adolescent twins are registered and their health and development monitored by school physicians and nurses. The tertiary prevention and health system (village, township, and county level) in Qingdao has provided assistance for the registries with high efficiency, coverage, and accuracy of information (Li et al., 2006). The twin registry and health-related surveys have been reviewed and approved by the

RECEIVED 31 August 2012; ACCEPTED 17 October 2012. First published online 27 November 2012.

ADDRESS FOR CORRESPONDENCE: Zengchang Pang, Qingdao Centers for Disease Control and Prevention, Shandong Road 175, Sifang District, Qingdao, 266033, P. R. China. Email: cdc-pang@126.com

QDCDC ethics committee and collaborative institutions' ethics committees.

## Anthropometric Measurements

Trained physicians or nurses explained the objectives of the survey and obtained consent from twins and their families. Height and weight were measured with participants wearing light clothes and without shoes. Waist circumference was measured at the midpoint between the rib cage and the iliac crest to the nearest 0.1 cm. Hip circumference was measured over the widest part of the gluteal region to the nearest centimeter. Body mass index (BMI) was defined as an individual's body weight in kilograms divided by the square of his/her height in meters ( $\text{kg/m}^2$ ). Blood pressure was taken by a standard procedure using mercurial table stand model sphygmomanometer. Systolic blood pressure was defined as Korotkoff of phase I (appearance of sound), and diastolic blood pressure was defined as Korotkoff phase V (disappearance of sound) (Ren et al., 2003). After undergoing the anthropometric measurements, adult twins, adolescent twins, and their parents were requested to fill in the health-related questionnaires, which included questions on substance use, mental health status of the twins, as well as socio-economic characteristics of the families.

## Laboratory Tests

After an overnight fast of 10–12 hours, twins were invited to the health screening centers of the local Center of Disease Control. Blood samples of approximately 10 mL were collected from the antecubital vein into a vacuum tube containing sodium fluoride. Centrifugation and separation of serum and plasma from blood cells were performed at the local centers within 30 minutes of collection. Fasting plasma glucose was determined using the glucose oxidase method. Total cholesterol (TC), triglycerides (TG), high-density lipoprotein cholesterol (HDL-C), and other biomedical parameters were measured using the Analyser Medical System (Olympus-AU 640 Automatic analyzers, Olympus Optical, Tokyo, Japan).

A standard 75 g oral glucose tolerance test was carried out in the adult twins. Participants with a history of diabetes or a fasting plasma glucose  $\geq 7.0$  mmol/L or a 2-hour plasma glucose  $\geq 11.1$  mmol/L, were classified as diabetic according to the World Health Organization criteria 1999 (Alberti & Zimmet, 1998). Another 5 mL venous blood sample was collected and stored for the DNA test.

## Zygosity Determination

For zygosity determination, first we identified potential monozygotic (MZ) and dizygotic (DZ) twins through sex and ABO blood types. Twins with opposite sex and/or different blood types were classified to DZ twins. Zygosity determination of twin pairs with same sex and blood types was classified by DNA testing using 16 short tandem repeat

**TABLE 1**

**Phenotypes of Twin Pairs for the Adolescents' Cohort**

Questionnaires
Twins
Education level and study performance
Relation with your co-twin
Tanner scale
Cohesion and conflict subscales of family adaptability
Life Events Scale
Depressive symptom
Hostility scale of SCL-90
Zuckerman sensation-seeking scale
Cigarette smoking
Alcohol consumption
Parents
Education and income level
Self-rating depression and anxiety scale
Cigarette smoking
Alcohol consumption
Anthropometric measurements for twins
Height, sitting height, weight, shoulder, waist, hip circumference, and body fat percentage
Blood pressure
Pulmonary function test (CO, FEV1, FVC)
Ophthalmology (optometry, diopter, funduscopy, slit lamp)
Biochemical measurements for twins
Cholesterol, HDL-cholesterol, triglycerides, glucose, BUN, and uric acid

Note: CO, carbon monoxide; FEV1, forced expiratory volume in 1 second; FVC, forced vital capacity; BUN, blood urea nitrogen.

markers (Becker et al., 1997; Jackson et al., 2001). Zygosity tests were conducted in the central laboratory of the Qingdao Blood Station.

## Development of the Adolescent Twin Cohort and the Research Focus

Six young twin cohorts (born 1990–1996) including a total of 600 pairs were established in 2006. The first phase of the young twin study was completed when these twins were 10–16 years old in 2006. The two follow-up surveys took place in 2007–2008 and in 2009 when twins were 11–18 and 13–19 years old, respectively. Follow-up rates were 98% in 2007–2008 and 95% in 2009. The questionnaires and anthropometric measurements were described in details as below (see Table 1).

Abnormal lipid metabolism contributes to atherosclerosis, cardiovascular diseases (Zhang et al., 2009), and diabetes (Mooradian, 2009) and lipid metabolism in adolescence is predictive for these conditions. Data from the QTR, including 314 same-sex twin pairs (MZ = 243, DZ = 71) aged 5–18 years, indicated that lipid levels were significantly different before and after puberty (Chen et al., 2009). Moderate and strong heritability estimates were observed for lipid phenotypes, such as TC, HDL-C, and low-density lipoprotein cholesterol (LDL-C), whereas TG was influenced mainly by shared and non-shared environmental factors. Total phenotypic variances decreased after puberty, mainly as a result of decrease of genetic variance, even though the common environmental variance increased. Shared environmental factors had an important effect on lipids during puberty,

suggesting family-based lifestyle intervention might be recommended in reducing cardiovascular risk factors. To what extent the variances change might be caused by sex and growth hormones or genetic-environmental interactions during puberty (Frindik, 1999; Rogol, 2002), needs to be investigated in a longitudinal study.

Hyperuricemia, defined as a serum uric acid concentration in excess of urate solubility (Fang & Alderman, 2000), increases the risk for diabetes and cardiovascular disease in adults. Accumulated evidence has demonstrated that hyperuricemia is associated with metabolic syndrome in child and adolescent Caucasians (Ford et al., 2007; Invitti et al., 2006; Lee et al., 2007; Pacifico et al., 2009; Tang et al., 2010), but not in their Chinese counterparts. Our data, including 521 twin pairs from the 2006 baseline survey ( $MZ = 274$ ,  $DZ = 247$ ), demonstrated that genetic factors play a vital role in uric acid metabolism. A sex-limitation model showed that additive genetic factors accounted for 75% of total variance, while non-shared environmental factors explained 22.5% (Duan et al., 2010). The combined effects of pubertal development, sex hormones, and obesity on uric acid levels need to be further investigated.

Skeletal age is usually employed to evaluate endocrine, genetic, and renal disorders and to monitor growth in childhood and adolescence (Poznanski, 1984; Tanner et al., 1975). The skeletal or bone age determination evaluates left hand and wrist from subjects at different stages of skeletal maturation using radiologic examinations (Greulich, 1959). Inconsistency between skeletal age and chronological age suggest metabolic disorders on the endocrinology and anthropology vary across the ethnic groups (Mora et al., 2001). During 2012–2013, a total of 190 pairs of adolescent twins from the QTR will have hand and wrist radiographs taken by experienced pediatric radiologists. The Standards of Skeletal Maturity of Hand and Wrist questionnaire (Version 2005) using measurements derived from Chinese children will be employed to determine maturation and subsequent evaluation of growth during puberty. The heritability of skeletal age and corresponding risk factors will be estimated in a sex limitation model. Whether the difference between skeletal age and chronological age is explained by genetic or environmental effects, such as sex hormone and lifestyle, will be assessed and compared with other ethnic groups.

#### Chinese–US Adolescents Twin Study of Substance Use

Cigarette smoking has been identified as the second leading risk factor for death from any cause worldwide (Gu et al., 2009). China is the world's largest producer and consumer of tobacco and bears a large proportion of deaths attributable to smoking worldwide. In 2010, a cross-sectional survey indicated that 28.1% of adults were current smokers, accounting for 301 million people in China (Li et al., 2011). Odds ratio (OR) for past 30-day smoking is significantly higher in adolescents in the United States than Chinese

counterparts in a multivariable adjusted model, with an OR of 0.80 (0.71–0.90; Unger et al., 2002). The association between access to cigarette and respondents' smoking were significantly greater in US adolescents. Family and peer influence play an important role on adolescent substance use (Trinidad et al., 2003; Unger et al., 2002; Zhu et al., 1992). Most psychological and genetic studies of adolescent substance use have been conducted within a single cultural context; however, it is not known whether the complex interactions among genetic risk/protective factors and social-environmental risk/protective factors differ across cultural contexts. Cross-cultural study could add to our knowledge about the socio-cultural conditions that increase or decrease the effects of risk and protective factors.

For this purpose, we initiated a cross-cultural study between China and the US adolescent twins based on 600 twin pairs (born between 1990 and 1996) in China and 600 twin pairs of the same age range selected from the twin study of University of Southern California in the United States. We collected psychosocial data to distinguish environmental risk and protective factors from heritable factors in order to identify the specific aspects of the shared and non-shared environments that are associated with substance use, and to investigate interactions and mediated pathways among these influences. Furthermore, the impact of parents and peers attitudes and behaviors on adolescents substance use will be investigated in this study.

Preliminary outcomes from the QTR cohort indicated that the estimates heritability for alcohol consumption is greater than for cigarette smoking (0.68 vs. 0.28), whereas environmental factors account for major components of total variances of psychological traits (depression, aggression, and hostility) among adolescents. Bivariate analysis showed that substance use and psychological measures were attributed to the combined effects of genetic and environmental factors (Unger et al., 2011). Family environment and peer behaviors were further investigated to explore the risk and protective factors for substance use in QTR. Family cohesion, family adaptability, parental monitoring, and academic score were associated with decreased risk for smoking, whereas negative events (e.g., family-related, school-related, and peer smoking) increased the odds for cigarette smoking in adolescents (Arpawong et al., 2010). Whether the association between risk and protective factors is culture specific warrants further investigations.

#### The Development of Adult Twin Survey and the Research Focus

A total of 695 pairs of adult twins ( $MZ = 405$ ,  $DZ = 290$ ) with a mean age of 37 years were recruited in this study. Anthropometric measurements of metabolic phenotypes were collected from these twins in 2001, and were followed up in 2004 and 2008. Genetic factors play vital roles in the sub-phenotypes of metabolic syndrome; however, no common

genetic mechanism interacts among the phenotypes based on the bivariate analyses (Duan et al., 2011). Multivariate analysis of lipid parameters explored the common genetic architecture in the development of these phenotypes. The findings revealed common genetic and environmental backgrounds for lipid phenotypes, that is, TC, HDL-C, and LDL-C. Modeling showed that physiologically distinct endophenotypes may follow different genetic regulations (Pang et al., 2010). To determine which loci affect lipid phenotypes in Chinese populations, genome-wide association analysis is warranted.

To our knowledge, we were the first to perform a genome-wide linkage scan on BMI using twins in the Chinese population. The strong linkage identified in Chinese subjects comparable with European studies suggests the existence of evolutionarily conserved genetic mechanisms for BMI, whereas the multiple suggestive loci could represent a genetic effect from gene-environment interaction as a result of population-specific environmental adaptation (Zhang et al., 2012a). Our genome-wide association analysis in DZ twins detected a suggestive association on chromosome 4 (rs17031508,  $p < 8, 34\text{E-}08$ ) located within a wide region of suggestive linkage. Our results provide some evidence for genetic linkage and associations with pulse pressure in the Chinese population. Further investigation is warranted to replicate the findings and to explore the susceptibility loci or genes for pulse pressure (Zhang et al., 2012b). Considering the limited sample size, validity of these loci requires additional replication studies to be conducted, particularly in the Chinese and other Asian populations. Moreover, the long-term effects of genetic and environmental risk factors for cardiovascular diseases will also be evaluated in a follow-up study.

### **The Development of Chinese–Finland Adolescents Twin Cohort Study and the Research Focus**

It is generally believed that genetic predisposition is important in the risk of disease, whereas lifestyle and environmental factors are assumed to trigger or modify the expression. It has been hypothesized that ethnic differences may result in varying susceptibility to diseases. For example, the risk of type 2 diabetes is 13.5 times greater in Asian children than in White children (Drake et al., 2002), while the incidence of type 1 diabetes is about 30 times higher in Finland than in China (LaPorte et al., 1985; Yang et al., 1998). Asthma is one of the most common atopic diseases in developed countries; however, the rates of asthma and allergy are relatively lower in Chinese children (Warner, 1999; Williams et al., 1999; Beasley, 1998). Genetic and environmental diversity, diet, substance use, and psychology might explain the regional variations. To what extent the differences in culture and lifestyle have contributed to the variations between Chinese and Finnish adolescents are less known. Twin studies should provide unique insights into how genetic and environmental factors influence obesity and other metabolic disorders

during urbanization and lifestyle transition in Chinese adolescents, and how comparable are the variance component estimates to those in Finnish counterparts.

We will collect data from 600 Chinese adolescent twins in a cohort established in 2006 when the pairs were 10–16 years old, and the data will be compared with 1,000 adolescent Finnish twin pairs at the same age. The specific aims of this study are: (1) to study interactions between genetic factors, social environment (e.g., parental and peer influences, stressful life events), and lifestyle (e.g., physical activity, cigarette smoking, and alcohol drinking) in the prediction of adolescent obesity and metabolic disorders in Chinese and Finnish adolescents; and (2) to assess mediated pathways by which heritable factors influence adolescents' selection of social environments, thereby influencing their exposure to social-environmental risk factors for obesity and metabolic disorders, making adolescents more susceptible or resilient to health consequences under different cultural contexts.

### **Development of Elderly Twin Cohort and the Research Focus**

Cross-sectional and longitudinal studies have shown that aging-related decline in physical performance and increasing functional impairment is associated with detrimental outcomes in the elderly, such as disability, decline in quality of life, and mortality. People over 60 years old account for 13.3% of the population in China. A collaborative study including 2,476 adults aged 50–74 years from the Qingdao China Ageing Study (Wu et al., 2012a, 2012b) and 5,278 twins aged 50–74 years from the Study of Middle-Aged Danish Twins and the Longitudinal Study of Aging Danish Twins, demonstrated significant gender and population differences in physical, cognitive performance and mental assessment (Wu et al. 2012a, 2012b). Age-related decline in general health was observed in both ethnic groups. However, genetic and environmental factors cannot be evaluated in the general population using the Chinese cohort.

To provide valuable information for decomposition of the genetic and environmental determinants in the observed differences in the two ethnic groups, we launched an elderly twin cohort in Qingdao. The protocol of elderly twin study is consistent with the Qingdao China Ageing Study (Wu et al. 2012a, 2012b). As shown in Table 2, physical performance, such as sit-to-stand test (lower limb strength measurement), handgrip strength, cognitive performance, and depression were evaluated in these elderly twins. The scale of the Minimal Cognitive state (Folstein et al., 1975) and the Zung Self-Rating Depression employed in the current study have been validated widely (Zung, 1965).

We plan to collect 300 pairs of twin aged 40 years and over to evaluate the genetic and environmental risk factors for aging-related diseases. The aims of the elderly twin study are: (1) to estimate the heritability of aging-related



**TABLE 2****Phenotypes of Twin Pairs for the Elder Cohort**

Questionnaires
Disease history
Socioeconomic status (education, income, and occupation)
Cognitive function
Self-Rated Health
Life Events Scale
Self-rating Depression Scale
Cigarette smoking and alcohol consumption
Physical activity
Anthropometric measurements
Height, weight, waist, hip circumference, and body fat percentage
Blood pressure
Sit-to-stand Test
Handgrip strength
Hearing Threshold
Pulmonary function test (CO, FEV1, FVC)
Ophthalmology
Biochemical measurements for twins
Cholesterol, HDL-cholesterol, triglycerides, glucose, BUN, and uric acid

phenotypes, and promote the well-being through environmental modifications; and (2) to explore the aging susceptible genes with the extensive genome-wide association studies.

Furthermore, we plan to conduct a longitudinal study in the adult and elderly twin cohorts. Meanwhile, together with the Qingdao Death Registry and Qingdao Health Record Registry, we monitor the incidence of diabetes, cancer, and other non-communicable diseases and lifestyle changes in Qingdao twins. An investigation of the joint effect of genetic and environmental risk factors on the increasing trends of metabolic disorders, mortality, and morbidity will be conducted in further investigations.

## Acknowledgments

The authors wish to thank the QDCDC twin project staff for assistance in data collection, and the twins and their families for their participation in this research. The QTR was supported by the China Medical Board, New York (CMB01-746). Additional support for expanding the twin registry was provided by the National Institutes of Health (P50 CA84735-01), Novo Nordisk Foundation Grant for Medical Research (2006), European Foundation for the Study of Diabetes (2007), and National Natural Science Foundation of China (30872170). FN also received financial support from European Foundation for the Study of Diabetes/Chinese Diabetes Society/Lilly Research Fellowships (2009).

## References

- Alberti, K. G., & Zimmet, P. Z. (1998). Definition, diagnosis and classification of diabetes mellitus and its complications. Part 1: Diagnosis and classification of diabetes mellitus provisional report of a WHO consultation. *Diabetic Medicine*, 15, 539–553.
- Arpawong, T. E., Sun, P., Chang, M. C., Gallaher, P., Pang, Z., Guo, Q., . . . Unger, J. (2010). Family and personal protective factors moderate the effects of adversity and negative disposition on smoking among Chinese adolescents. *Substance Use & Misuse*, 45, 1367–1389.
- Beasley, R. (1998). Worldwide variation in prevalence of symptoms of asthma, allergic rhinoconjunctivitis, and atopic eczema: ISAAC. The international study of asthma and allergies in childhood (ISAAC) steering committee. *Lancet*, 351, 1225–1232.
- Becker, A., Busjahn, A., Faulhaber, H. D., Bähring, S., Robertson, J., Schuster, H., & Luft, F. C. (1997). Twin zygosity: Automated determination with microsatellites. *The Journal of Reproductive Medicine*, 42, 260–266.
- Chen, T. J., Ji, C. Y., & Hu, Y. H. (2009). Genetic and environmental influences on serum lipids and the effects of puberty: A Chinese twin study. *Acta Paediatrica*, 98, 1029–1036.
- Chan, J. C., Malik, V., Jia, W., Kadowaki, T., Yajnik, C. S., Yoon, K. H., & Hu, F. B. (2009). Diabetes in Asia: Epidemiology, risk factors, and pathophysiology. *The Journal of the American Medical Association*, 301, 2129–2140.
- China's major health challenge: Control of chronic diseases. (2011). *Lancet*, 378, 457.
- Drake, A. J., Smith, A., Betts, P. R., Crowne, E. C., & Shield, J. P. (2002). Type 2 diabetes in obese white children. *Archives of Disease in Childhood*, 86, 207–208.
- Duan, H., Pang, Z., Zhang, D., Li, S., Kruse, T. A., Kyvik, K. O., . . . Tan, Q. (2011). Genetic and environmental dissections of sub-phenotypes of metabolic syndrome in the Chinese population: A twin-based heritability study. *Obesity Facts*, 4, 99–104.
- Duan, H. P., Pang, Z. C., Zhang, D., Wang, S. J., Zhai, Y. M., & Zhang, D. F. (2010). Heritability of serum uric acid in adolescent twins. *Acta Academiae Medicinae Qingdao Universitatis*, 45, 415–418.
- Fang, J., & Alderman, M. H. (2000). Serum uric acid and cardiovascular mortality the NHANES I epidemiologic follow-up study, 1971–1992. National Health and Nutrition Examination Survey. *The Journal of the American Medical Association*, 283, 2404–2410.
- Folstein, M. F., Folstein, S. E., & McHugh, P. R. (1975). 'Mini-mental state': A practical method for grading the cognitive state of patients for the clinician. *Journal of Psychiatric Research*, 12, 189–198.
- Ford, E. S., Li, C., Cook, S., & Choi, H. K. (2007). Serum concentrations of uric acid and the metabolic syndrome among US children and adolescents. *Circulation*, 115, 2526–2532.
- Frindik, J. P., Kemp, S. F., & Sy, J. P. (1999). Effects of recombinant human growth hormone on height and skeletal maturation in growth hormone-deficient children with and without severe pretreatment bone age delay. *Hormone Research*, 51, 15–19.
- Greulich, W. W., & Pyle, S. L. (1959). *Radiographic skeletal development of the hand and wrist* (2nd ed.). Stanford, CA: Stanford University Press.
- Gu, D., Kelly, T. N., Wu, X., Chen, J., Samet, J. M., Huang, J. F., . . . He, J. (2009). Mortality attributable to smoking in China. *The New England Journal of Medicine*, 360, 150–159.
- Invitti, C., Maffei, C., Gilardini, L., Pontiggia, B., Mazzilli, G., Girola, A., . . . Viberti, G. C. (2006). Metabolic syndrome in

- obese Caucasian children: Prevalence using WHO-derived criteria and association with nontraditional cardiovascular risk factors. *International Journal of Obesity*, 30, 627–633.
- Jackson, R. W., Snieder, H., Davis, H., & Treiber, F. A. (2001). Determination of twin zygosity: A comparison of DNA with various questionnaire indices. *Twin Research*, 4, 12–18.
- LaPorte, R. E., Tajima, N., Akerblom, H. K., Berlin, N., Brosseau, J., Christy, M., . . . Neil, A. (1985). Geographic differences in the risk of insulin-dependent diabetes mellitus: The importance of registries. *Diabetes Care*, 8 (Suppl. 1), 101–107.
- Lee, M. S., Wahlqvist, M. L., Yu, H. L., & Pan, W. H. (2007). Hyperuricemia and metabolic syndrome in Taiwanese children. *Asia Pacific Journal of Clinical Nutrition*, 16 (Suppl. 2), 594–600.
- Li, L., Gao, W., Lv, J., Cao, W., Zhan, S., Yang, H., & Hu, Y. (2006). Current status of the Chinese national twin registry. *Twin Research and Human Genetics*, 9, 747–752.
- Li, Q., Hsia, J., & Yang, G. (2011). Prevalence of smoking in China in 2010. *The New England Journal of Medicine*, 364, 2469–2470.
- Mooradian, A. D. (2009). Dyslipidemia in type 2 diabetes mellitus. *Nature Clinical Practice Endocrinology & Metabolism*, 5, 150–159.
- Mora, S., Boechat, M. I., Pietka, E., Huang, H. K., & Gilsanz, V. (2001). Skeletal age determinations in children of European and African descent: Applicability of the Greulich and Pyle standards. *Pediatric Research*, 50, 624–628.
- Pacifico, L., Cantisani, V., Anania, C., Bonaiuto, E., Martino, F., Pascone, R., & Chiesa, C. (2009). Serum uric acid and its association with metabolic syndrome and carotid atherosclerosis in obese children. *European Journal of Endocrinology/European Federation of Endocrine Societies*, 160, 45–52.
- Pang, Z., Bo, T., Wang, S., Peng, X., Zhang, Y., & Zhai, Y. (2003). Analysis on recruitment of monodizymus in Qingdao. *China Journal of Public Health*, 19, 66–67.
- Pang, Z., Ning, F., Unger, J., Johnson, C. A., Wang, S., Guo, Q., . . . Lee, L. (2006). The Qingdao twin registry: A focus on chronic disease research. *Twin Research and Human Genetics*, 9, 758–762.
- Pang, Z., Zhang, D., Li, S., Duan, H., Hjelmberg, J., Kruse, T. A., . . . Tan, Q. (2010). Multivariate modelling of endophenotypes associated with the metabolic syndrome in Chinese twins. *Diabetologia*, 53, 2554–2561.
- Poznanski, A. K. (1984). *The hand in radiologic diagnosis with gamuts and pattern profiles* (2nd ed.). Philadelphia: WB Saunders.
- Ren, T., Wu, D., Hu, Y., Cao, W., Zhan, S., Lv, J., . . . Li, L. (2003). The heritability analysis of the metabolic syndrome related characters in twins. *Chinese Journal of Prevention and Control of Chronic Diseases*, 11, 13–15.
- Rogol, A. D., Roemmich, J. N., & Clark, P. A. (2002). Growth at puberty. *Journal of Adolescent Health*, 31 (6 Suppl.), 192–200.
- Tang, L., Kubota, M., Nagai, A., Mamemoto, K., & Tokuda, M. (2010). Hyperuricemia in obese children and adolescents: The relationship with metabolic syndrome. *Pediatric Reports*, 2, e12.
- Tanner, J. M., Whitehouse, R. H., & Marshall, W. A. (1975). *Assessment of skeletal maturity and prediction of adult height (TW2 method)*. New York: Academic Press.
- Trinidad, D. R., Chou, C. P., Unger, J. B., Johnson, C. A., & Li, Y. (2003). Family harmony as a protective factor against adolescent tobacco and alcohol use in Wuhan, China. *Substance use & Misuse*, 38, 1159–1171.
- Unger, J. B., Lessov-Schlaggar, C. N., Pang, Z., Guo, Q., Ning, F., Gallaher, P., . . . Anderson Johnson, C. (2011). Heritability of smoking, alcohol use, and psychological characteristics among adolescent twins in Qingdao, China. *Asia-Pacific Journal of Public Health*, 23, 568–580.
- Unger, J. B., Yan, L., Shakib, S., Rohrbach, L. A., Chen, X., Qian, G., . . . Anderson Johnson, C. (2002). Peer influences and access to cigarettes as correlates of adolescent smoking: A cross-cultural comparison of Wuhan, China, and California. *Preventive Medicine*, 34, 476–484.
- Wang, L., Kong, L., Wu, F., Bai, Y., & Burton, R. (2005). Preventing chronic diseases in China. *Lancet*, 366, 1821–1824.
- Warner, J. O. (1999). Worldwide variations in the prevalence of atopic symptoms: What does it all mean? *Thorax*, 54, S46–S51.
- Williams, H., Robertson, C., Stewart, A., Ait-Khaled, N., Anabwani, G., Anderson, R., . . . Weiland, M. D. Jr. (1999). Worldwide variations in the prevalence of symptoms of atopic eczema in the international study of asthma and allergies in childhood. *The Journal of Allergy and Clinical Immunology*, 103, 125–138.
- World Health Organization (WHO). 1994. Prevention of diabetes mellitus (Report of a WHO study group). *World Health Organization Technical Report Series*, 844, 1–100.
- Wu, Y., Pang, Z., Zhang, D., Jiang, W., Wang, S., Li, S., . . . Tan, Q. (2012a). A cross-sectional analysis of age and sex patterns in grip strength, tooth loss, near vision and hearing levels in Chinese aged 50–74 years. *Archives of Gerontology and Geriatrics*, 54, e213–e220.
- Wu, Y., Zhang, D., Pang, Z., Oksuzyan, A., Jiang, W., Wang, S., . . . Tan, Q. (2012b). Gender-specific patterns in age-related decline in general health among Danish and Chinese: A cross-national comparative study. *Geriatrics & Gerontology International*, 12, 431–439.
- Yang, Z., Wang, K., Li, T., Sun, W., Li, Y., Chang, Y.F., . . . LaPorte, R. E. (1998). Childhood diabetes in China. Enormous variation by place and ethnic group. *Diabetes Care*, 21, 525–529.
- Zhang, D. F., Pang, Z., Li, S., Thomassen, M., Wang, S., Jiang, W., . . . Tan, Q. (2012a). High-resolution genome-wide linkage mapping identifies susceptibility loci for BMI in the Chinese population. *Obesity (Silver Spring)*, 20, 830–833.
- Zhang, D. F., Pang, Z., Li, S., Jiang, W., Wang, S., Thomassen, M., . . . Tan, Q. (2012b). Genome-wide linkage and association scans for pulse pressure in Chinese twins. *Hypertension Research*. Advance online publication. doi:10.1038/hr.2012.90.

- Zhang, L., Qiao, Q., Tuomilehto, J., Hammar, N., Ruotolo, G., Stehouwer, C. D. A., . . . for DECODE/DECODA Study Group. (2009). The impact of dyslipidaemia on cardiovascular mortality in individuals without a prior history of diabetes in the DECODE Study. *Atherosclerosis*, 206, 298–302.
- Zhu, B. P., Liu, M., Wang, S. Q., He, G. Q., Chen, D. H., Shi, J. H., . . . Shang, J. Z. (1992). Cigarette smoking among junior high school students in Beijing, China, 1988. *International Journal of Epidemiology*, 21, 854–861.
- Zung, W. W. (1965). A self-rating depression scale. *Archives of General Psychiatry*, 12, 63–70.
-